

DOI: 10.1002/adsc.201301109

Palladium-Catalyzed Decarbonylative Dehydration of Fatty Acids for the Production of Linear Alpha Olefins

Yiyang Liu,^a Kelly E. Kim,^a Myles B. Herbert,^a Alexey Fedorov,^a
Robert H. Grubbs,^{a,*} and Brian M. Stoltz^{a,*}^a The Warren and Katharine Schlinger Laboratory for Chemistry and Chemical Engineering, Division of Chemistry and Chemical Engineering, California Institute of Technology, 1200 E. California Blvd, MC 101-20, Pasadena, CA 91125, USA
Fax: (+1)-626-395-8436; e-mail: rhg@caltech.edu or stoltz@caltech.edu

Received: December 9, 2013; Published online: January 13, 2014

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/adsc.201301109>.


Abstract: A highly efficient palladium-catalyzed decarbonylative dehydration reaction of carboxylic acids is reported. This method transforms abundant and renewable even-numbered natural fatty acids into valuable and expensive odd-numbered alpha olefins. Additionally, the chemistry displays a high functional group tolerance. The process employs a low loading of palladium catalyst and proceeds under solvent-free and relatively mild conditions.

Keywords: decarbonylative dehydration; fatty acids; large-scale process; linear alpha olefins; palladium; renewable feedstocks; Z-selective metathesis

Linear alpha olefins represent an important class of industrial chemicals with a wide range of applications. They are used as co-monomers for ethylene polymerization^[1] as well as precursors to plasticizers, lubricants, and surfactants.^[2] Currently, these olefins are mainly produced by oligomerization of ethylene^[3] which, in turn, is derived from petroleum. As the world's oil reserves continue to diminish, the development of renewable feedstocks for the production of alpha olefins will become increasingly important. One obvious choice is ethylene from biomass-derived ethanol.^[4] A potentially more direct method is the decarbonylative dehydration of long-chain fatty acids. The latter route is particularly attractive because fatty acids are inexpensive and readily available starting materials derived from many natural sources. Since natural fatty acids contain an even number of carbon atoms, their corresponding alpha olefins will be odd-numbered after decarbonylative dehydration. Moreover, conventional ethylene oligomerization processes deliver only even-numbered alpha olefins,^[3] and odd-

numbered olefins are largely inaccessible and prohibitively expensive.^[5] These odd-numbered olefins are valuable building blocks in the synthesis of various fine chemicals such as lepidopteran insect pheromones, but are currently far too costly to be practical.^[6] Therefore, the development of an efficient and economic process for fatty acid decarbonylative dehydration is highly desirable.

Many strategies to convert fatty acids to alpha olefins have been pursued. Lead tetraacetate-mediated oxidative decarboxylation is a classical method.^[7] Alternative protocols that avoid stoichiometric toxic reagents have also been developed, such as Kolbe electrolysis^[8] and silver-catalyzed oxidative decarboxylation.^[9] However, these reactions proceed through highly reactive radical intermediates, and thus suffer from low yields due to many side reactions. A more recent approach entails transition metal-catalyzed decarbonylative dehydration of fatty acids. Various transition metals including rhodium,^[10] iridium,^[11] palladium,^[12] and iron^[13] have been shown to catalyze decarbonylative dehydration reactions. To date, palladium has demonstrated the highest activity, and catalyst loadings as low as 0.01 mol% have been reported independently by Miller^[12a] and Kraus^[12b] (Scheme 1A). Unfortunately, their methods require very high temperatures (230–250 °C). In addition, it is necessary to distill the olefin product from the reaction mixture as soon as it is formed in order to prevent double bond isomerization, and therefore only volatile olefins can be produced this way. Decarbonylation processes under milder conditions have been developed independently by Gooßen^[12c] and Scott^[12d] (Scheme 1B). Although their reactions proceed at 110 °C, much higher palladium catalyst loading (3 mol%) and an expensive, high-boiling-point solvent (DMPU) are required. We envisioned that by judicious choice of the ligand set for palladium and other parameters, the most advantageous aspects of these two systems could



Scheme 1. Palladium-catalyzed decarbonylative dehydration. **(A)** High temperature processes (Miller, Kraus). **(B)** Low temperature processes (Gooßen, Scott). **(C)** This research.


be combined. Herein we report a palladium-catalyzed decarbonylative dehydration using low catalyst loading under relatively mild and solvent-free conditions to produce alpha olefins in good yield and high selectivity (Scheme 1C).


At the outset of our study, we examined the palladium-catalyzed decarbonylative dehydration of stearic acid **1a** in neat acetic anhydride as the dehydrating agent (Table 1). A preliminary survey of phosphine ligands revealed Xantphos to be an optimal and unique ligand for the transformation (entries 1–4). It is believed that acetic anhydride converts the stearic acid into stearic anhydride *in situ*, which then undergoes oxidative addition by Pd(0) to initiate the catalytic cycle.^[12] Considering that acetic anhydride itself could compete with stearic anhydride for oxidative addition at the metal center, we sought to simplify the system by using pre-formed stearic anhydride **1a'** alone. To our surprise, the decarbonylation of neat stearic anhydride with the same catalyst was exceedingly slow (only 12% yield, 120 TON, in 2 h; entry 5). Comparing the two systems, we found that the former had one equivalent of acid in it, while the latter was acid-free. Thus, we posited that acid might play a role in promoting reactivity. Consequently we added 1 mol% isophthalic acid to the system, and the yield rose to 22% (entry 6). When the ligand-to-metal ratio was reduced from 4:1 to 1.2:1, the yield dramatically increased to 92%, but the selectivity dropped to 31% (entry 7). Since it has been shown previously that triphenylphosphine can inhibit olefin isomerization,^[12a] we used PdCl₂(PPh₃)₂ instead of PdCl₂(nbd) (nbd = norbornadiene), and we were delighted to observe an increase in selectivity to 54% with negligible erosion

in yield (90%; entry 8). Finally, we examined a number of protic additives (entries 9–12), and found that (*t*-Bu)₄biphenol gave the optimal overall performance to furnish the highest yield of alpha olefin (entry 12).




One major limitation of using pre-formed stearic anhydride as the substrate is that only half of the molecule can be converted to the olefin while the other half is wasted as stearic acid, so the maximum theoretical yield based on the acid is 50%. Control experiments also revealed that the build-up of acid in the reaction mixture was responsible for olefin isomerization and erosion of alpha selectivity. We envisioned that portionwise addition of acetic anhydride to the reaction mixture and distillation of acetic acid *in situ* could transform stearic acid back into its anhydride, thereby reducing acid concentration and increasing olefin yield and selectivity (Table 2). When two portions of acetic anhydride were added to the reaction mixture with immediate distillation of acetic acid, once every 1.5 h, the olefin product was obtained in 69% yield based on stearic acid and 62% selectivity (entry 1). Furthermore, when three portions of acetic anhydride were added, once every hour, the olefin product was obtained in 67% yield and 86% alpha selectivity (entry 2). Finally, when six portions of acetic anhydride were added, once every half hour, the olefin product was obtained in 67% isolated yield and 89% alpha selectivity (entry 3). The selectivity trend clearly shows that the more frequently acetic anhydride is added to dehydrate stearic acid, the higher is the alpha selectivity. This is a remarkable result in that it is possible to maintain high selectivity without

Table 1. Effects of catalyst, ligand, and additive.^[a]

A. 

B. 

| Entry | Rxn | Pd cat. | Ligand (mol%) | Additive | Yield [%] ^[b] | Alpha [%] ^[b] | Y x A [%] ^[c] |
|-------|-----|--|------------------------|------------------------------|--------------------------|--------------------------|--------------------------|
| 1 | A | PdCl ₂ (nbd) | PPh ₃ (0.8) | -- | 0 | -- | 0 |
| 2 | A | PdCl ₂ (nbd) | dppp (0.4) | -- | 0 | -- | 0 |
| 3 | A | PdCl ₂ (nbd) | DPEphos (0.4) | -- | 43 | 59 | 25 |
| 4 | A | PdCl ₂ (nbd) | Xantphos (0.4) | -- | 60 | 55 | 33 |
| 5 | B | PdCl ₂ (nbd) | Xantphos (0.4) | -- | 12 | 100 | 12 |
| 6 | B | PdCl ₂ (nbd) | Xantphos (0.4) | isophthalic acid | 22 | 96 | 21 |
| 7 | B | PdCl ₂ (nbd) | Xantphos (0.12) | isophthalic acid | 92 | 31 | 29 |
| 8 | B | PdCl ₂ (PPh ₃) ₂ | Xantphos (0.12) | isophthalic acid | 90 | 54 | 49 |
| 9 | B | PdCl ₂ (PPh ₃) ₂ | Xantphos (0.12) | p-TsOH·H ₂ O | 86 | 5 | 4 |
| 10 | B | PdCl ₂ (PPh ₃) ₂ | Xantphos (0.12) | salicylamide | 60 | 90 | 54 |
| 11 | B | PdCl ₂ (PPh ₃) ₂ | Xantphos (0.12) | 2,2'-biphenol | 59 | 91 | 54 |
| 12 | B | PdCl ₂ (PPh ₃) ₂ | Xantphos (0.12) | (t-Bu) ₄ biphenol | 84 | 70 | 59 |



^[a] Conditions: **A**) 1 equiv. **1a** (5 mmol), 2 equiv. Ac₂O; **B**) 1 equiv **1a'** (5 mmol).

^[b] Determined by ¹H NMR with methyl benzoate as internal standard. Alpha = **2a**/(**2a**+**3a**).

^[c] Y x A = Yield x Alpha.

having to distill the olefin product out of the reaction mixture as it forms.

With the optimized conditions in hand, we explored the decarbonylative dehydration of a variety of carboxylic acid substrates, as shown in Table 3. Common saturated fatty acids with carbon numbers from 12 to 18 all provided the corresponding olefin in good yield and high alpha selectivity (entries 1–4). In particular, volatile olefins were formed with exceptionally high selectivities (entries 3 and 4). Terminally functionalized fatty acids were also competent substrates, and functional groups such as esters, chlorides, imides, silyl ethers, ketones, terminal olefins, and substituted aromatics were all well tolerated (entries 5–13). Notably, allylbenzene derivative **2k** was formed with 91% alpha selectivity (entry 11), which is impressive con-

sidering the significant thermodynamic driving force for isomerization into conjugation with the aromatic ring. Carboxylic acids with α- or β-substituents were considerably less reactive (entries 14–16). Nevertheless, catalyst turnovers up to 400 could be achieved for these challenging substrates. Compared to previous reports by Miller^[12a] and Kraus,^[12b] our reaction exhibits a much broader scope, does not require distillation of the olefin products to maintain high selectivity, and is compatible with various heteroatom-containing functional groups.

Since this process requires no solvent and low catalyst loading, it can be readily scaled up (Scheme 2). In a laboratory setting, a 100-mmol scale (28.4 g stearic acid or 23.0 g 10-acetoxydecanoic acid) decarbonylative dehydration was easily carried out in a 100-mL

Table 2. Portionwise addition of acetic anhydride.^[a]

| | | | |
|--|--|--------------------------|--------------------------|
| $ \begin{array}{c} \text{C}_{15}\text{H}_{31}\text{CH}_2\text{CH}_2\text{COOH} \\ \textbf{1a} \\ + \\ \text{Ac}_2\text{O (portionwise)} \end{array} \xrightarrow[\text{neat, 132 } ^\circ\text{C, 3 h}]{\text{PdCl}_2(\text{PPh}_3)_2 \text{ (0.05 mol\%)} \\ \text{Xantphos (0.06 mol\%)} \\ \text{(t-Bu)}_4\text{biphenol (0.5 mol\%)}} \begin{array}{c} \text{C}_{15}\text{H}_{31}\text{CH=CH}_2 \\ \textbf{2a} \\ + \\ \text{internal olefins} \\ \textbf{3a} \end{array} $ | | | |
| Entry | Equiv. of Ac ₂ O | Yield [%] ^[b] | Alpha [%] ^[b] |
| 1 | 1+0.5 (once every 1.5 hours) | 69 | 62 |
| 2 | 1+0.5+0.25 (once every hour) | 67 | 86 |
| 3 | 1+0.14+0.12+0.1+0.09+0.08 (once every half hour) | 68 (67) | 89 |

^[a] 20 mmol **1a**.

^[b] Determined by ¹H NMR (isolated yield in parentheses).

round-bottom flask. Compared with small-scale reactions, the large-scale ones afforded products in similar yields and slightly higher alpha selectivity. When the olefin is sufficiently volatile, it is distilled out together with the acetic acid (Table 3, entries 3–6, 11, and 13). Although distillation of olefin is not necessary in order to maintain high selectivity (e.g., Scheme 2A), it is convenient to do so for volatile olefins (Scheme 2B).


The olefin products thus obtained are important building blocks in chemical synthesis. For example, 8-nonenyl acetate (**2f**) is a precursor to insect pheromones **4a** (oriental fruit moth pheromone)^[14] and **4b** (figure-of-eight moth pheromone).^[15] Aided by *Z*-selective cross-metathesis catalyst **5**, olefin **2f** reacts with 1-pentene or 1-hexene to afford pheromones **4a** or **4b** in 58% and 48% yields, respectively, and >98% *Z*-selectivity (Table 4).^[6b]

In summary, we have developed a highly efficient palladium-catalyzed decarbonylative dehydration process that converts carboxylic acids (e.g., fatty acids) to linear alpha olefins in good yield and with high selectivity. The reaction requires low palladium catalyst loading and proceeds under solvent-free and relatively mild conditions. *In situ* distillation of the olefin product is not necessary, and a wide range of functionalized and unfunctionalized carboxylic acids can be transformed into their corresponding olefins. Process development for large-scale production is underway and will be reported in due course.

Experimental Section

General Procedure

A 15-mL round-bottom flask was charged with PdCl₂(PPh₃)₂ (0.01 mmol, 0.05 mol%), Xantphos (0.012 mmol, 0.06 mol%), (t-Bu)₄biphenol (0.1 mmol, 0.5 mol%), and fatty acid substrate (20 mmol, 1 equiv.). The flask was equipped with a distillation head and a 25-mL round-bottom receiving flask. The closed system was connected to a vacuum manifold, equipped with a needle valve and a digital vacuum gauge. The system was evacuated and refilled with N₂ (×3), and the first portion of acetic anhydride (20 mmol, 1 equiv.) was added *via* syringe through the septum that seals the top of the distillation head. The flask was lowered into an oil bath at 20 °C and gradually heated to 132 °C in 23 min. When the oil bath temperature had risen to 122 °C, the needle valve was closed, switched to vacuum, and the needle valve carefully and slowly opened to allow distillation of acetic acid into a receiving flask, which was cooled to –78 °C. When the oil bath temperature reached 130 °C, time was recorded as t=0. After distillation ceased (about t=3 min), the needle valve was opened fully and a vacuum of 1–5 mmHg was drawn. At t=30 min, the system was refilled with N₂, and the second portion of acetic anhydride (2.8 mmol, 0.14 equiv.) was added *via* syringe. The system was then gradually (t=35 min) resubjected to



Scheme 2. Large-scale decarbonylative dehydration of stearic acid and 10-acetoxydecanoic acid.

Table 3. Substrate scope study.^[a]

| $ \begin{array}{c} \text{R}-\text{CH}_2-\text{CH}_2-\text{COOH} \\ \mathbf{1} \\ + \text{Ac}_2\text{O (6 portions)} \end{array} \xrightarrow[\text{neat, 132 } ^\circ\text{C, 3 h}]{\text{PdCl}_2(\text{PPh}_3)_2 \text{ (0.05 mol\%)} \\ \text{Xantphos (0.06 mol\%)} \\ \text{(t-Bu)}_4\text{biphenol (0.5 mol\%)} \\ \text{1–5 mmHg distillation} \\ \text{– CO, – AcOH}} \begin{array}{c} \text{R}-\text{CH}=\text{CH}_2 \\ \mathbf{2} \\ + \\ \text{internal olefins} \\ \mathbf{3} \end{array} $ | | | | | |
|--|-----------|---------|--------------------------|---------------------------|--|
| Entry | Substrate | Product | Yield [%] ^[b] | TON | Alpha [%] ^[c] |
| 1 | | | 67 | 1340 | 89 |
| 2 | | | 41 | 820 | 97 |
| 3 | | | 65 | 1300 | 99 |
| 4 | | | 73 | 1460 | 99 |
| 5 ^[d,e] | | | 63 | 1260 | 98 |
| 6 ^[d] | | | 67 | 1340 | 96 |
| 7 ^[d] | | | 60 | 1200 | 89 |
| 8 | | | 75 | 1500 | 86 |
| 9 | | | 76 | 1520 | 83 |
| 10 | | | 64 | 1280 | 80 |
| 11 | | | 80 | 1600 | 91 |
| 12 | | | 49 | 980 | 88 |
| 13 | | | 59 | 1180 | 87 |
| 14 ^[f] | | | 20 80 ^[h] | 400 320 ^[h] | ... ^[g] ... ^[h] |
| 15 | | | 19 | 380 | ... ^[g] |
| 16 ^[i] | | | 71 | 71 | ... ^[i] |

- [a] Conditions: 20 mmol **1**, 6 portions of Ac₂O, 1 + 0.14 + 0.12 + 0.10 + 0.09 + 0.08 equiv., added every 30 min.
 [b] Isolated yield (column chromatography).
 [c] Determined by ¹H NMR.
 [d] Purified by distillation.
 [e] 18.5 mmol **1e**.
 [f] PdCl₂(nbd) (0.05 mol%), PPh₃ (0.05 mol%), Xantphos (0.06 mol%), 1.5 h, 3 portions of Ac₂O (1 + 0.15 + 0.10 equiv.).
 [g] Single isomer observed.
 [h] PdCl₂(PPh₃)₂ (0.25 mol%), Xantphos (0.30 mol%), (*t*-Bu)₄biphenol (1 mol%), **2n:3n** = 49:51.
 [i] 2-Methyldecanoic anhydride (10 mmol), no Ac₂O, PdCl₂(nbd) (1 mol%), Xantphos (1.1 mol%), salicylamide (2 mol%), 160 °C, 10 mmHg distillation, 10 h, **3p:2p** = 73:27.

Table 4. Synthesis of pheromones **4a** and **4b** from olefin **2f**.

$$\text{R}-\text{CH}=\text{CH}_2 + \text{CH}_2=\text{CH}(\text{CH}_2)_6\text{OAc} \xrightarrow[\text{THF, 35 } ^\circ\text{C, 2 h}]{\text{5 (0.5 mol\%)}} \text{R}-\text{CH}=\text{CH}(\text{CH}_2)_6\text{OAc}$$

5 =

| Product | R | Yield [%] ^[a] | Z [%] ^[b] |
|-----------|-------------------------------|--------------------------|----------------------|
| 4a | C ₃ H ₇ | 58 | >98 |
| 4b | C ₄ H ₉ | 48 | >98 |

[a] 4.8 mmol **2f**.

[b] Determined by ¹H NMR.

a vacuum of 1–5 mmHg. Acetic anhydride was added as follows (0.12, 0.10, 0.09, 0.08 equiv.) in the same manner every 30 min. The reaction was stopped at *t* = 3 h and allowed to cool to ambient temperature. The residual reaction mixture was purified by flash chromatography. If it contained solids, it was suction-filtered first and the solids washed with hexanes, and the filtrate was concentrated and purified by chromatography. In cases where the product was distilled together with acetic acid, the distillate was added dropwise to a saturated NaHCO₃ solution, stirred for 30 min, and the resulting mixture was extracted with dichloromethane (30 mL × 3). The combined extracts were dried over Na₂SO₄, filtered and concentrated. The crude product was then subjected to flash chromatography or distillation to afford the olefin in pure form.

Acknowledgements

The authors wish to thank the Resnick Sustainability Institute at Caltech (graduate fellowship to Y.L.), BP (postdoctoral fellowship to A.F. under the XC2 program), NIH-NIGMS (R01GM080269, B.M.S.), NIH (NIH 5R01GM031332-27, R.H.G.), the Gordon and Betty Moore Foundation, the Cal-

tech Center for Catalysis and Chemical Synthesis, and Caltech for financial support. Matera, Inc. is thanked for the generous donation of ruthenium catalysts. Dr. David VanderVelde is acknowledged for NMR spectroscopy assistance. Dr. Mona Shahgholi and Naseem Torian are acknowledged for high-resolution mass spectrometry assistance. Dr. Scott Virgil is acknowledged for helpful discussions.

References

- [1] S. D. Ittel, L. K. Johnson, M. Brookhart, *Chem. Rev.* **2000**, *100*, 1169–1203.
 - [2] a) D. M. Marquis, S. H. Sharman, R. House, W. A. Sweeney, *J. Am. Oil Chem. Soc.* **1966**, *43*, 607–614; b) R. Franke, D. Selent, A. Börner, *Chem. Rev.* **2012**, *112*, 5675–5732.
 - [3] a) A. M. Al-Jarallah, J. A. Anabtawi, M. A. B. Siddiqui, A. M. Aitani, A. W. Al-Sa'doun, *Catal. Today* **1992**, *14*, 1–121; b) B. L. Small, M. Brookhart, *J. Am. Chem. Soc.* **1998**, *120*, 7143–7144.
 - [4] J. Ouyang, F. Kong, G. Su, Y. Hu, Q. Song, *Catal. Lett.* **2009**, *132*, 64–74.
 - [5] Prices of acids and olefins (Sigma–Aldrich, 10/5/2013):
- | acid | \$/kg | olefin | \$/kg |
|-----------------------------|-------|----------------------------------|-----------|
| decanoic (C ₁₀) | 16.28 | 1-nonene (C ₉) | 12,080.00 |
| | | 1-decene (C ₁₀) | 55.87 |
| lauric (C ₁₂) | 14.20 | 1-undecene (C ₁₁) | 7,240.00 |
| | | 1-dodecene (C ₁₂) | 187.60 |
| myristic (C ₁₄) | 11.76 | 1-tridecene (C ₁₃) | 23,800.00 |
| | | 1-tetradecene (C ₁₄) | 47.74 |
| palmitic (C ₁₆) | 7.68 | 1-pentadecene (C ₁₅) | 24,722.58 |
| | | 1-hexadecene (C ₁₆) | 2,758.62 |
| stearic (C ₁₈) | 13.20 | 1-heptadecene (C ₁₇) | 69,500.00 |
- [6] a) M. B. Herbert, V. M. Marx, R. L. Pederson, R. H. Grubbs, *Angew. Chem.* **2013**, *125*, 328–332; *Angew. Chem. Int. Ed.* **2013**, *52*, 310–314; b) L. E. Rosebrugh, M. B. Herbert, V. M. Marx, B. K. Keitz, R. H. Grubbs, *J. Am. Chem. Soc.* **2013**, *135*, 1276–1279.
 - [7] J. D. Bacha, J. K. Kochi, *Tetrahedron* **1968**, *24*, 2215–2226.

- [8] H. Kolbe, *Justus Liebigs Ann. Chem.* **1849**, 69, 257–294.
- [9] a) J. M. Anderson, J. K. Kochi, *J. Org. Chem.* **1970**, 35, 986–989; b) F. van der Klis, M. H. van den Hoorn, R. Blaauw, J. van Haveren, D. S. van Es, *Eur. J. Lipid Sci. Technol.* **2011**, 113, 562–571.
- [10] T. A. Foglia, P. A. Barr, *J. Am. Oil Chem. Soc.* **1976**, 53, 737–741.
- [11] S. Maetani, T. Fukuyama, N. Suzuki, D. Ishihara, I. Ryu, *Organometallics* **2011**, 30, 1389–1394.
- [12] a) J. A. Miller, J. A. Nelson, M. P. Byrne, *J. Org. Chem.* **1993**, 58, 18–20; b) G. A. Kraus, S. Riley, *Synthesis* **2012**, 44, 3003–3005; c) L. J. Gooßen, N. Rodríguez, *Chem. Commun.* **2004**, 724–725; d) J. Le Nôtre, E. L. Scott, M. C. R. Franssen, J. P. M. Sanders, *Tetrahedron Lett.* **2010**, 51, 3712–3715; e) M. O. Miranda, A. Pietrangelo, M. A. Hillmyer, W. B. Tolman, *Green Chem.* **2012**, 14, 490–494.
- [13] S. Maetani, T. Fukuyama, N. Suzuki, D. Ishihara, I. Ryu, *Chem. Commun.* **2012**, 48, 2552–2554.
- [14] T. C. Baker, R. T. Cardé, J. R. Miller, *J. Chem. Ecol.* **1980**, 6, 749–758.
- [15] M. Subchev, M. Toth, D. Wu, L. Stanimirova, T. Toshova, Z. Karpati, *J. Appl. Entomol.* **2000**, 124, 197–199.