


Palladium-Catalyzed Enantioselective C_{sp^3} – C_{sp^3} Cross-Coupling for the Synthesis of (Poly)fluorinated Chiral Building BlocksYanhui Lu,^{†,‡} Elizabeth L. Goldstein,[‡] and Brian M. Stoltz^{*,‡,§}[†]Graduate School of Engineering, Nagoya University, Furo-cho, Chikusa, Nagoya 464-8603, Japan[‡]The Warren and Katharine Schlinger Laboratory for Chemistry and Chemical Engineering, Division of Chemistry and Chemical Engineering, California Institute of Technology, MC 101-20, Pasadena, California 91125, United States

S Supporting Information



ABSTRACT: A general method for the enantioselective synthesis of carbo- and heterocyclic carbonyl compounds bearing fluorinated α -tetrasubstituted stereocenters using palladium-catalyzed decarboxylative allylic alkylation is described. The stereoselective C_{sp^3} – C_{sp^3} cross-coupling reaction delivers five- and six-membered ketone and lactam products bearing (poly)fluorinated tetrasubstituted chiral centers in high yields and enantioselectivities. These fluorinated, stereochemically rich building blocks hold potential value in medicinal chemistry and are prepared using an orthogonal and enantioselective approach into such chiral moieties compared to traditional approaches, often without the use of electrophilic fluorinating reagents.

Organofluorine compounds often play a critical role in the lead optimization phase of drug discovery, due to their impact on various physicochemical properties such as absorption, distribution, metabolic stability, and excretion. Consequently, more than 20% of marketed pharmaceuticals contain C–F motifs, despite the fact that organofluorinated compounds are exceedingly rare in nature.¹ Recently, molecules with tetrasubstituted stereocenters have attracted the interest of medicinal chemists aiming to incorporate three-dimensionality and added novelty.² Importantly, there are many successful marketed pharmaceuticals bearing fluorinated tetrasubstituted stereocenters (1–3, Figure 1). For these




Figure 1. Marketed active pharmaceutical ingredients bearing fluorinated tetrasubstituted stereocenters.

reasons, there has been renewed interest in the synthesis of fluorinated tetrasubstituted stereocenters for use in drug discovery and, in particular, access to new classes of fluorinated analogs. Therefore, we believe that a general method for the construction of fluorine-containing tetrasubstituted stereo-

centers will be of particular interest to chemists in the area of drug discovery and development.


Methods to construct fluorine-containing α -tetrasubstituted ketones have been the subject of intense investigation over the past decade. The most prevalent strategy for fluorine incorporation is intermolecular catalytic asymmetric electrophilic fluorination (or trifluoromethylation) of enolates (Scheme 1A).³ Despite their potential utility in organic synthesis, the relatively low abundance of cheap, commercially available electrophilic fluorinating and trifluoromethylating reagents prohibits their widespread usage. As far back as 2005, the Stoltz and Nakamura groups independently reported the intramolecular asymmetric allylic alkylation of prochiral enolates derived from the decarboxylation of 1,3-dicarbonyl substrates (Scheme 1B).^{4,5} Using this strategy, several optically active α -fluoro α -tetrasubstituted cyclic carbonyl derivatives have been synthesized in high yield and enantioselectivity.

While stereogenic C–F moieties have been previously investigated, the compatibility of fluoroalkyl groups in palladium-catalyzed asymmetric allylic alkylation has remained unknown until recently. In 2011, Shibata and co-workers reported the first example of the construction of trifluoromethyl-bearing quaternary centers by intramolecular decarboxylative allylic alkylation of α -trifluoromethyl β -ketoesters (Scheme 1C).⁶ Unfortunately, attempts to render their reaction enantioselective were unsuccessful. Due to our interest in the field of asymmetric allylic alkylation, we endeavored to build on these previous reports and investigate a number of fluoroalkyl and fluoroallyl derivatives in


Received: July 26, 2018

Scheme 1. Asymmetric Construction of Fluorine-Containing α -Tetrasubstituted Ketones


A: Intermolecular fluorinations (or trifluoromethylation)³




B: Intramolecular allylic alkylation to introduce C–F tetrasubstituted stereocenters⁵



C: Intramolecular allylic alkylation to introduce C–CF₃ quaternary stereocenters⁶



D: This research




asymmetric allylic alkylation reactions. Herein, we report the first general method for the construction of carbo- and heterocyclic carbonyl derivatives bearing α -fluoro-, α -fluoroalkyl-, or α -(2-fluoro)allyl substituents using palladium-catalyzed enantioselective decarboxylative allylic alkylation (Scheme 1D).

Importantly, with this strategy, a number of fluorinated alkyl and allyl groups are introduced into the substrate via standard 1,3-dicarbonyl chemistry (thermal, acidic or basic conditions) to produce racemic mixtures of compounds that serve as substrates for the mild and neutral asymmetric allylic alkylation reaction. In some cases, these fluorinated substrates are synthesized without the use of electrophilic fluorinating reagents. Furthermore, this allows for the nonasymmetric formation of the C–F or C–CF₃ bonds, which are significantly more developed than their asymmetric equivalents. For example, 1,1,1-trifluoropropyl groups can be installed using standard β -keto ester alkylation conditions utilizing 1,1,1-trifluoropropyl iodide and base in moderate yields (Scheme 2A). The synthesis of 1,1,1-trifluoroethyl substituted β -keto esters proceeded smoothly with the use of 2,2,2-trifluoroethyl (mesityl)iodonium trifluoromethanesulfonate⁷ (available in 2 steps from commercial materials) in the presence of LiHMDS (Scheme 2B). During the preparation of this manuscript, a report using 2,2,2-trifluoroethyl(mesityl) iodonium trifluoromethanesulfonate for the alkylation of 1,3-dicarbonyls was disclosed using similar conditions.⁸


In addition to α -fluoroalkyl groups, a number of 2-fluoro allyl substrates were prepared without the use of electrophilic fluorinating reagents. Starting from commercially available methyl 2-fluoroacrylate, reduction of the ester to the alcohol occurred, followed by treatment with 1,1'-carbonyldiimidazole resulting in the formation of an acylating reagent (Scheme 2C). This reagent could then be used as previously reported⁹ to form a β -keto ester (Scheme 2D), which can be subsequently alkylated or fluorinated.^{4,5} Additionally, using known chemistry, α -fluoro β -keto esters can be synthesized using Selectfluor^{sd} and α -trifluoromethyl β -keto esters can be

Scheme 2. Synthesis of Fluorinated β -Ketoesters


A: Synthesis of α -trifluoropropyl β -ketoester




B: Synthesis of α -trifluoroethyl β -ketoester



C: Synthesis of 2-fluoroallyl acyl imidazole



D: Synthesis of 2-fluoroallyl β -ketoester




synthesized using Umemoto's Reagent,⁶ both of which are commercially available.

Initial reaction optimization started with trifluoroethyl substituted β -ketoester **4a** using catalytic Pd₂(dba)₃ at 23 °C in diethyl ether in the presence of a chiral PHOX ligand toward the synthesis of ketone **5a** (Table 1).¹⁰ Employing the

Table 1. Optimization of Conditions for Enantioselective Palladium-Catalyzed Allylic Alkylation^a

entry	ligand	solvent	yield (%)	ee (%) ^b
1	(S)- <i>t</i> -BuPHOX	Et ₂ O	88	85
2	(S)-(CF ₃) ₃ - <i>t</i> -BuPHOX	Et ₂ O	99	90
3	(S)-(CF ₃) ₃ - <i>t</i> -BuPHOX	THF	93	86
4	(S)-(CF ₃) ₃ - <i>t</i> -BuPHOX	TBME	95	90
5	(S)-(CF ₃) ₃ - <i>t</i> -BuPHOX	toluene	99	90


^aConditions: β -ketoester **4a** (0.1 mmol), Pd₂(dba)₃ (5.0 mol %), ligand (12.5 mol %), solvent (3 mL). ^bDetermined by analytical chiral SFC.



classic (*S*)-*t*-BuPHOX ligand, the desired product was formed in 88% yield and 85% ee (entry 1). By switching to the electron-deficient (*S*)-(CF₃)₃-*t*-BuPHOX ligand, the desired product was furnished in an improved 99% yield and 90% ee (entry 2). Solvent effects were not very significant (entries 3–5); however, THF gave a decreased ee of 86% (entry 3), while the less polar TBME and nonpolar toluene performed similarly to diethyl ether. Based on these results, we determined that using Pd₂(dba)₃ (5.0 mol %) with (*S*)-(CF₃)₃-*t*-BuPHOX in toluene (0.033 M) at room temperature proved optimal.¹¹

Subsequently, we explored the substrate scope of the enantioselective allylic alkylation of fluorine-containing 1,3-dicarbonyl compounds (Scheme 3). We found that our

Scheme 3. Substrate Scope of Fluorine-Containing compounds in Enantioselective Allylic Alkylation^a



^aUnless otherwise noted, all reported yields are isolated yields. Enantiomeric excess (ee) was determined by chiral SFC. Standard conditions: β -ketoester **5** (0.1 mmol), Pd₂(dba)₃ (5 mol %), (*S*)-(CF₃)₃-*t*-BuPHOX (12.5 mol %), toluene (3 mL), 23 °C, 24 h. ^bReaction performed at 40 °C. ^cReaction performed in the presence of Pd₂(pmdba)₃ instead of Pd₂(dba)₃. ^dReaction performed at 60 °C. ^eReaction performed at 23 °C for 70 h.

reaction was tolerant of a variety of α -fluoro-, α -fluoroalkyl-, and α -fluoroallyl substituents to deliver five- and six-membered ketone and lactam products bearing fluorinated tetrasubstituted stereocenters in high yields and enantioselectivities. Trifluoropropyl substituted **4b** exhibited similar enantioinduction as **4a** to furnish **5b** in 92% ee and an extremely high yield. α -Fluoro tetrasubstituted compounds, which are usually introduced by direct fluorination with fluorine reagents and chiral catalysts,³ were prepared in a very efficient manner with high enantioselectivity (**5c**, **5d**), even in the presence of a chloroallyl substituent (**5d**). Surprisingly, 2-fluoroallyl groups

survived the palladium-catalyzed allylic alkylation even at elevated temperatures (40 °C),¹² albeit with slightly decreased enantioselectivity (**5e**). Recently, Shibata and co-workers described that enantioenriched indanone α -trifluoromethyl β -ketoesters lost their optical activity under the palladium-catalyzed allylic alkylation reaction conditions in the presence of achiral ligands to deliver a racemic α -quaternary ketone, and when they tried to render the transformation enantioselective, they were unsuccessful.⁶ However, we were pleased to see that α -trifluoromethyl substituted tetralone substrate **4f** reacted to furnish **5f** with a moderate level of enantioselectivity. Generally, five-membered cyclic β -ketoesters have performed worse than the corresponding six-membered ring substrates, often providing the α -tetrasubstituted ketone products in comparatively low ee.^{4k} Under these conditions, alkylation of the five-membered indanone substrates **4g** and **4h** occurred with levels of enantioinduction similar to those observed for the tetralone substrates, with only a slightly diminished 87% ee for trifluoropropyl-substituted indanone **5h**. Indanone substrates bearing a 2-fluoroallyl substituent proceeded in high yield, but with only moderate enantioselectivity, to form products **5i** and **5j**, following the trend of the 2-fluoroallyl tetralone substrates. Gratifyingly, lactam substrates were also well tolerated in the reaction. Trifluoropropyl-substituted *N*-benzoyl δ -valerolactam (**5k**) was obtained in 94% yield and 89% ee. Surprisingly, in contrast to the negative influence of the 2-fluoroallyl substituent on substrates **5e**, **5i**, and **5j**, the fluorine on the allyl group of *N*-benzoyl δ -valerolactam **4l** enhanced the enantioselectivity, providing **5l** in 97% ee. Additionally, trifluoropropyl-substituted *N*-benzyloxy glutarimide **5m** was furnished in 89% ee with high yield. Finally, *N*-benzoyl pyrrolidinone **5n** was obtained in diminished yield and ee.

In conclusion, we have developed a general method to construct fluorine-containing tetrasubstituted stereocenters by enantioselective palladium-catalyzed decarboxylative allylic alkylation. A strategy was adopted with the preintroduction of fluorine on racemic substrates, which could be used as an orthogonal approach to the traditional fluorination and trifluoromethylation strategies. The reaction manifold demonstrated significant substitution tolerance to furnish a wide range of five- and six-membered ketone and lactam products bearing fluorinated tetrasubstituted stereocenters in high yields and enantioselectivities. Furthermore, we provide the first examples demonstrating that 2-fluoroallyl substituents can survive in the presence of certain palladium sources and deliver related fluoroalkylated products in elevated enantiopurity.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.8b02369.

Experimental procedures and compound characterization (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: stoltz@caltech.edu.

ORCID

Brian M. Stoltz: 0000-0001-9837-1528

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

The NIH-NIGMS (R01GM080269) and Caltech are thanked for support of our research program. E.L.G. recognizes the NSF for a predoctoral research fellowship (No. DGE-1745301). Y.L. thanks the Program of Leading Graduate Schools: IGER Program in Green Natural Sciences (MEXT) at Nagoya University for financial support.

■ REFERENCES

- (1) For reviews on fluorine in medicinal chemistry, see: (a) Böhm, H.-J.; Banner, D.; Bendels, S.; Kansy, M.; Kuhn, B.; Müller, K.; Obst-Sander, U.; Stahl, M. *ChemBioChem* **2004**, *5*, 637. (b) Isanbor, C.; O'Hagan, D. *J. Fluorine Chem.* **2006**, *127*, 303. (c) Kirk, K. L. *J. Fluorine Chem.* **2006**, *127*, 1013. (d) Müller, K.; Faeh, C.; Diederich, F. *Science* **2007**, *317*, 1881. (e) Kirk, K. L. *Org. Process Res. Dev.* **2008**, *12*, 305. (f) O'Hagan, D. *Chem. Soc. Rev.* **2008**, *37*, 308. (g) Purser, S.; Moore, P. R.; Swallow, S.; Gouverneur, V. *Chem. Soc. Rev.* **2008**, *37*, 320. (h) Hagmann, W. K. *J. Med. Chem.* **2008**, *51*, 4359. (i) Qiu, X.-L.; Xu, X.-H.; Qing, F.-L. *Tetrahedron* **2010**, *66*, 789. (j) Hunter, L. *Beilstein J. Org. Chem.* **2010**, *6*, 38. (k) Vulpatti, A.; Dalvit, C. *Drug Discovery Today* **2012**, *17*, 890. (l) Wang, J.; Sánchez-Roselló, M.; Aceña, J. L.; del Pozo, C.; Sorochinsky, A. E.; Fustero, S.; Soloshonok, V. A.; Liu, H. *Chem. Rev.* **2014**, *114*, 2432. (m) Gillis, E. P.; Eastman, K. J.; Hill, M. D.; Donnelly, D. J.; Meanwell, N. A. *J. Med. Chem.* **2015**, *58*, 8315.
- (2) Lovering, F.; Bikker, J.; Humblet, C. *J. Med. Chem.* **2009**, *52*, 6752.
- (3) For reviews on fluorination and trifluoromethylation to construct α -tetrasubstituted stereocenters on ketones, see: (a) Bobbio, C.; Gouverneur, V. *Org. Biomol. Chem.* **2006**, *4*, 2065. (b) Prakash, G. K. S.; Beier, P. *Angew. Chem., Int. Ed.* **2006**, *45*, 2172. (c) Pihko, P. M. *Angew. Chem., Int. Ed.* **2006**, *45*, 544. (d) Hamashima, Y.; Sodeoka, M. *Synlett* **2006**, *2006*, 1467. (e) Shibata, N.; Ishimaru, T.; Nakamura, S.; Toru, T. *J. Fluorine Chem.* **2007**, *128*, 469. (f) Ma, J.-A.; Cahard, D. *Chem. Rev.* **2008**, *108*, PR1. (g) Shibata, N.; Mizuta, S.; Kawai, H. *Tetrahedron: Asymmetry* **2008**, *19*, 2633. (h) Brunet, V. A.; O'Hagan, D. *Angew. Chem., Int. Ed.* **2008**, *47*, 1179. (i) Ueda, M.; Kano, T.; Maruoka, K. *Org. Biomol. Chem.* **2009**, *7*, 2005. (j) Lectard, S.; Hamashima, Y.; Sodeoka, M. *Adv. Synth. Catal.* **2010**, *352*, 2708. (k) Cahard, D.; Xu, X.; Couve-Bonnaire, S.; Pannecoucke, X. *Chem. Soc. Rev.* **2010**, *39*, 558. (l) Zheng, Y.; Ma, J.-A. *Adv. Synth. Catal.* **2010**, *352*, 2745. (m) Shibata, N.; Matsnev, A.; Cahard, D. *Beilstein J. Org. Chem.* **2010**, *6*, 65. (n) Furuya, T.; Kamlet, A. S.; Ritter, T. *Nature* **2011**, *473*, 470. (o) Valero, G.; Companyó, X.; Rios, R. *Chem. - Eur. J.* **2011**, *17*, 2018. (p) Hollingworth, C.; Gouverneur, V. *Chem. Commun.* **2012**, *48*, 2929. (q) Macé, Y.; Magnier, E. *Eur. J. Org. Chem.* **2012**, *2012*, 2479. (r) Liang, T.; Neumann, C. N.; Ritter, T. *Angew. Chem., Int. Ed.* **2013**, *52*, 8214. (s) Yang, X.; Wu, T.; Phipps, R. J.; Toste, F. D. *Chem. Rev.* **2015**, *115*, 826.
- (4) For accounts and reviews on palladium-catalyzed allylic alkylation chemistry, see: (a) Trost, B. M.; Van Vranken, D. L. *Chem. Rev.* **1996**, *96*, 395. (b) Trost, B. M. *Acc. Chem. Res.* **1996**, *29*, 355. (c) Pfaltz, A.; Lautens, M. In *Comprehensive Asymmetric Catalysis* III, Vol. 2; Jacobsen, E. N., Pfaltz, A., Yamamoto, H., Eds.; Springer: New York, 1999; pp 833–884. (d) Helmchen, G. *J. Organomet. Chem.* **1999**, *576*, 203. (e) Trost, B. M.; Lee, C. In *Catalytic Asymmetric Synthesis*, 2nd ed.; Ojima, I., Ed.; Wiley: New York, 2000; pp 593–649. (f) Trost, B. M. *Chem. Pharm. Bull.* **2002**, *50*, 1. (g) Graening, T.; Schmalz, H.-G. *Angew. Chem.* **2003**, *115*, 2684; *Angew. Chem., Int. Ed.* **2003**, *42*, 2580. (h) Trost, B. M. *J. Org. Chem.* **2004**, *69*, 5813. (i) Lu, Z.; Ma, S. *Angew. Chem.* **2008**, *120*, 264; *Angew. Chem., Int. Ed.* **2008**, *47*, 258. (j) Behenna, D. C.; Mohr, J. T.; Sherden, N. H.; Marinescu, S. C.; Harned, A. M.; Tani, K.; Seto, M.; Ma, S.; Novák, Z.; Krout, M. R.; McFadden, R. M.; Roizen, J. L.; Enquist, J. A., Jr.; White, D. E.; Levine, S. R.; Petrova, K. V.; Iwashita, A.; Virgil, S. C.; Stoltz, B. M. *Chem. - Eur. J.* **2011**, *17*, 14199. (k) Liu, Y.; Han, S.-J.; Liu, W.-B.; Stoltz, B. M. *Acc. Chem. Res.* **2015**, *48*, 740. (5) (a) Mohr, J. T.; Behenna, D. C.; Harned, A. M.; Stoltz, B. M. *Angew. Chem., Int. Ed.* **2005**, *44*, 6924. (b) Nakamura, M.; Hajra, A.; Endo, K.; Nakamura, E. *Angew. Chem.* **2005**, *117*, 7414; *Angew. Chem., Int. Ed.* **2005**, *44*, 7248. (c) Behenna, D. C.; Liu, Y.; Yurino, T.; Kim, J.; White, D. E.; Virgil, S. C.; Stoltz, B. M. *Nat. Chem.* **2012**, *4*, 130. (d) Mohr, J. T.; Nishimata, T.; Behenna, D. C.; Stoltz, B. M. *J. Am. Chem. Soc.* **2006**, *128*, 11348. (6) Shibata, N.; Suzuki, S.; Furukawa, T.; Kawai, H.; Tokunaga, E.; Yuan, Z.; Cahard, D. *Adv. Synth. Catal.* **2011**, *353*, 2037. (7) Tolnai, G. L.; Székely, A.; Makó, Z.; Gáti, T.; Daru, J.; Bihari, T.; Stirling, A.; Novák, Z. *Chem. Commun.* **2015**, *51*, 4488. (8) Zhao, C.-L.; Yang, J.; Han, Z.-Z.; Zhang, C.-P. *J. Fluorine Chem.* **2017**, *204*, 23. (9) Trost, B. M.; Xu, J. *J. Org. Chem.* **2007**, *72*, 9372. (10) Absolute configuration of **5c** was determined by comparison of the optical rotation of the same compound to the known literature value; see: Nakamura, M.; Hajra, A.; Endo, K.; Nakamura, E. *Angew. Chem.* **2005**, *117*, 7414; *Angew. Chem., Int. Ed.* **2005**, *44*, 7248. The absolute configuration of all other products generated herein was assigned by analogy to the absolute configuration of **5c**. For full details, see the [Supporting Information](#). (11) With the consideration of heating requirements for some less reactive substrates, toluene was assigned as the best solvent. (12) A defluorinated side product was obtained in the presence of Pd(PPh₃)₄ for preparing racemic standards. Therefore, racemic samples were prepared in the presence of Pd₂(dba)₃ (or Pd₂(pmdba)₃) and achiral Gly-PHOX for fluoro-allyl products. For full details, see the [Supporting Information](#).