

Pd-Catalyzed Enantioselective Aerobic Oxidation of Secondary Alcohols: Applications to the Total Synthesis of Alkaloids

Shyam Krishnan, Jeffrey T. Bagdanoff, David C. Ebner, Yeeman K. Ramtohl,
Uttam K. Tambar, and Brian M. Stoltz*

*The Arnold and Mabel Beckman Laboratories of Chemical Synthesis, Division of Chemistry and
Chemical Engineering, California Institute of Technology, 1200 East California Boulevard,
MC 164-30, Pasadena, California 91125*

Received June 20, 2008; E-mail: stoltz@caltech.edu

Abstract: Enantioselective syntheses of the alkaloids (–)-aurantioclavine, (+)-amurensinine, (–)-lobeline, and (–)- and (+)-sedamine are described. The syntheses demonstrate the effectiveness of the Pd-catalyzed asymmetric oxidation of secondary alcohols in diverse contexts and the ability of this methodology to set the absolute configuration of multiple stereocenters in a single operation. The utility of an aryne C–C insertion reaction in accessing complex polycyclic frameworks is also described.

Introduction

Chiral amines are a key functional group in numerous biologically active natural products and synthetic drugs. The enantioselective synthesis of amines has been extensively investigated and continues to be a field of intense interest.¹ Chiral amines can be accessed from the corresponding alcohols using several possible approaches. This transformation could, in principle, be achieved via functional group interconversion of a hydroxyl to an amine functionality at a stereogenic carbon (Scheme 1a, path A).² Alternatively, desymmetrization of a *meso*-substrate bearing multiple pro-stereogenic centers with alcohol and amine functional groups could deliver the corre-

sponding enantioenriched product (Scheme 1a, path B).³ In addition, kinetic resolution of a racemic, diastereomerically pure substrate with stereocenters bearing hydroxyl and amine functionalities provides a route to chiral amino alcohols (Scheme 1a, path C). Finally, diastereoselective installation of a stereocenter bearing an amine group can be accomplished from a substrate that already features a hydroxyl-bearing stereocenter (Scheme 1a, path D). In this article, we present syntheses of natural products bearing chiral amines or chiral amino alcohols employing each of the above strategies.


We sought to apply the Pd-catalyzed aerobic oxidative kinetic resolution⁴ (Scheme 1b) developed in our laboratory toward the synthesis of alkaloids. This transformation enables the catalytic enantioselective synthesis of chiral alcohols, with benzylic alcohols being particularly excellent substrates.⁵ We present here syntheses of the natural products (–)-aurantioclavine ((–)-1), (+)-amurensinine ((+)-2), (–)-lobeline ((–)-3), (–)-sedamine ((–)-4), and (+)-sedamine (Figure 1) by kinetic resolution or desymmetrization of benzylic alcohols as the key steps.

(–)-Aurantoclavine ((–)-1) is an ergot alkaloid that was first isolated from *Penicillium aurantiovirens*.⁶ The alkaloid possesses a 3,4,5,6-tetrahydro-6-(2-methyl-1-propenyl)azepino[5,4,3-


- (1) For recent reviews on the enantioselective synthesis of amines, see: (a) Moody, C. J. *Angew. Chem., Int. Ed.* **2007**, *46*, 9148. (b) Ouellet, S. G.; Walji, A. M.; MacMillan, D. W. C. *Acc. Chem. Res.* **2007**, *40*, 1327. (c) Skucas, E.; Ngai, M.-Y.; Komanduri, V.; Krische, M. J. *Acc. Chem. Res.* **2007**, *40*, 1394. (d) Masson, G.; Housseman, C.; Zhu, J. *Angew. Chem., Int. Ed.* **2007**, *46*, 4614. (e) Li, G.; Saibau, K.; Timmons, C. *Eur. J. Org. Chem.* **2007**, *17*, 2745. (f) Bräse, S.; Baumann, T.; Dahmen, S.; Vogt, H. *Chem. Commun.* **2007**, 1881. (g) Maruoka, K.; Ooi, T.; Kano, T. *Chem. Commun.* **2007**, 1487. (h) Ellman, J. A. *Pure Appl. Chem.* **2003**, *75*, 39. (i) Roesky, P. W.; Mueller, T. E. *Angew. Chem., Int. Ed.* **2003**, *42*, 2708. (j) Enders, D.; Reinhold, U. *Tetrahedron: Asymmetry* **1997**, *8*, 1895.
- (2) For selected examples of synthesis of amines directly from alcohols, their corresponding esters, carbonates, sulfonate esters, or alkyl halides, see: (a) Kan, T.; Fukuyama, T. *Chem. Commun.* **2004**, 353. (b) Fujita, K.-i.; Fujii, T.; Yamaguchi, R. *Org. Lett.* **2004**, *6*, 3525. (c) Evans, P. A.; Robinson, J. E.; Nelson, J. D. *J. Am. Chem. Soc.* **1999**, *121*, 6761. (d) Scriven, E. F. V.; Turnbull, K. *Chem. Rev.* **1988**, *88*, 297. (e) Kolasa, T.; Miller, M. J. *J. Org. Chem.* **1987**, *52*, 4978. (f) Bestmann, H. J.; Wölfel, G. *Chem. Ber.* **1984**, *117*, 1250. (g) Trost, B. M.; Keinan, E. *J. Org. Chem.* **1979**, *44*, 3451. (h) Nordlander, J. E.; Catalane, D. B.; Eberlein, T. H.; Farkas, L. V.; Howe, R. S.; Stevens, R. M.; Tripoulas, N. A.; Stansfield, R. E.; Cox, J. L. *Tetrahedron Lett.* **1978**, *50*, 4987. (i) Jonczyk, A.; Ochal, Z.; Makosza, M. *Synthesis* **1978**, 882. (j) Zwierzak, A.; Podstawczynska, I. *Angew. Chem., Int. Ed. Engl.* **1977**, *16*, 702. (k) Hendrickson, J. B.; Joffe, I. *J. Am. Chem. Soc.* **1973**, *95*, 4083. (l) Mitsunobu, O.; Wada, M.; Sano, T. *J. Am. Chem. Soc.* **1972**, *94*, 679. (m) Gibson, M. S.; Bradshaw, R. W. *Angew. Chem., Int. Ed. Engl.* **1968**, *7*, 919.

- (3) For recent reviews of asymmetric desymmetrization, see: (a) Rendler, S.; Ostreich, M. *Angew. Chem., Int. Ed.* **2008**, *47*, 248. (b) Rovis, T. Recent Advances in Catalytic Asymmetric Desymmetrization Reactions. In *New Frontiers in Asymmetric Catalysis*; Mikami, K., Lautens, M., Eds.; John Wiley & Sons, Inc.: Hoboken, NJ, 2007; pp 275–311. (c) Atodiresei, I.; Schiffrers, I.; Bolm, C. *Chem. Rev.* **2007**, *107*, 5683. (d) Schneider, C. *Synthesis* **2006**, 23, 3919. (e) Garcia-Urdiales, E.; Alfonso, I.; Gotor, V. *Chem. Rev.* **2005**, *105*, 313.
- (4) (a) Ferreira, E. M.; Stoltz, B. M. *J. Am. Chem. Soc.* **2001**, *123*, 7725. (b) Jensen, D. R.; Pugsley, J. S.; Sigman, M. S. *J. Am. Chem. Soc.* **2001**, *123*, 7475.
- (5) For an experimentally derived model for stereoselectivity in the oxidative kinetic resolution of benzylic alcohols catalyzed by Pd-(sparteine)Cl₂, see: Trend, R. M.; Stoltz, B. M. *J. Am. Chem. Soc.* **2004**, *126*, 4482.
- (6) Kozlovskii, A. G.; Solov'eva, T. F.; Sahkarovskii, V. G.; Adanin, V. M. *Dokl. Akad. Nauk SSSR* **1981**, *260*, 230.

Scheme 1



Scheme 2



cd]indole tricyclic ring system bearing a single stereocenter. We were particularly interested in aurantioclavine as an intermediate en route to the complex polycyclic alkaloids of the communesin family⁷ (Scheme 2). While syntheses of racemic aurantioclavine⁸ have been reported in the literature, an asymmetric synthesis of aurantioclavine has not been reported, to the best of our knowledge. Our objective was to develop an enantioselective synthesis of (-)-1 as a means toward an asymmetric synthesis of members of the communesin family.⁹

(-)-Amurensinine ((-)-2) belongs to a family of alkaloids, the isopavines (Figure 2), originally isolated from Papaveraceae plants.¹⁰ The isopavines display biological activity relevant to neurological disorders such as Parkinson's disease, Down's syndrome, Alzheimer's disease, amyotrophic lateral sclerosis, and Huntington's chorea. Despite the potential medicinal applications of the isopavines and their analogues,¹¹ relatively few total syntheses of these natural products have been reported.¹² The majority of these syntheses involve intramolecular acid-promoted cyclization to form the azabicyclo[3.2.2]nonane core




Figure 1. Alkaloids with nitrogen-bearing stereocenters.




Figure 2. Isopavine natural products.

of the isopavines. We envisioned a catalytic enantioselective approach to rapidly access the isopavine core in a modular fashion.¹³

(-)-Lobeline ((-)-3) is a primary alkaloid constituent of *Lobelia inflata*, a plant commonly known as “Indian tobacco” because it was previously used by native North Americans as a tobacco substitute.^{14,15} A respiratory stimulant, the plant’s crude extracts have been widely used for the treatment of respiratory illnesses, including asthma, bronchitis, pneumonia, and whooping cough. (-)-3 mildly mimics the effect of nicotine, is an antagonist of nicotine acetylcholine receptors, and has thus been applied as a smoking cessation agent.¹⁶ (-)-3 also inhibits


the neurochemical and behavioral effects of methamphetamine and is an inhibitor of dopamine and vesicular monoamine transporter function. (-)-3 is thus a promising lead in the development of treatments for methamphetamine abuse.¹⁷ While it may be isolated from its natural source and crystallized as a single isomer in salt form, in the solution state the free base of (-)-3 is known to exist in equilibrium with its epimer at C(3).¹⁸ Due to this known equilibrium, early enantioselective syntheses¹⁹ by Marazano and Lebreton report the synthesis of (-)-3 as a mixture with its C(3) epimer. Our goal was to access the alkaloid 3 in diastereomerically pure form, and we believed that the application of the Pd-catalyzed enantioselective oxidation would allow efficient access to (-)-3 from a symmetric *meso*-intermediate. We also believed that access to either enantiomer of sedamine (4),^{20,21} a piperidine derivative found in various *Sedum* species, would be possible through the application of a Pd-catalyzed oxidative kinetic resolution.

Results and Discussion

Retrosynthesis of (-)-Aurantioclavine. Our retrosynthetic analysis for (-)-aurantioclavine ((-)-1) is depicted in Scheme


- (7) (a) Numata, A.; Takahashi, C.; Ito, Y.; Takada, T.; Kawai, K.; Usami, Y.; Matsumura, E.; Imachi, M.; Ito, T.; Hasegawa, T. *Tetrahedron Lett.* **1993**, *34*, 2355. (b) Jadulco, R.; Edrada, R. A.; Ebel, R.; Berg, A.; Schaumann, K.; Wray, V.; Steube, K.; Proksch, P. *J. Nat. Prod.* **2004**, *67*, 78. (c) Hayashi, H.; Matsumoto, H.; Akiyama, K. *Biosci. Biotechnol. Biochem.* **2004**, *68*, 753. (d) Dalsgaard, P. W.; Blunt, J. W.; Munro, M. H. G.; Frisvad, J. C.; Christophersen, C. *J. Nat. Prod.* **2005**, *68*, 258.
- (8) For total syntheses of aurantioclavine, see: (a) Somei, M.; Yamada, F. *Heterocycles* **2007**, *74*, 943. (b) Yamada, F.; Makita, Y.; Suzuki, T.; Somei, M. *Chem. Pharm. Bull.* **1985**, *33*, 2162. (c) Hegedus, L. S.; Toro, J. L.; Miles, W. H.; Harrington, P. J. *J. Org. Chem.* **1987**, *52*, 3319.
- (9) (a) May, J. A.; Stoltz, B. M. *Tetrahedron* **2006**, *62*, 5262. (b) May, J. A.; Zeidan, R. K.; Stoltz, B. M. *Tetrahedron Lett.* **2003**, *44*, 1203.
- (10) (a) Boit, H. G.; Flentje, H. *Naturwissenschaften* **1960**, *47*, 180. (b) Gözler, B.; Lantz, M. S.; Shamma, M. *J. Nat. Prod.* **1983**, *46*, 293.
- (11) For discussions of the biological activity of the isopavine alkaloids, see: (a) Weber, E.; Keana, J.; Barmettler, P. PCT Int. Appl. 106019w, 1991; *Chem. Abstr.* 1991, *115*, 106019w. (b) Childers, W. E.; Abou-Gharbia, M. A. U.S. Patent 4,940,789, 1990; *Chem. Abstr.* 1990, *113*, 191190w.
- (12) For selected syntheses of isopavine alkaloids, see: (a) Gözler, B. Pavine and Isopavine Alkaloids. In *The Alkaloids*; Brossi, A., Ed.; Academic Press: New York, 1987; Vol. 31, pp 343–356. (b) Meyers, A. I.; Dickman, D. A.; Boes, M. *Tetrahedron* **1987**, *43*, 5095. (c) Gottlieb, L.; Meyers, A. I. *J. Org. Chem.* **1990**, *55*, 5659. (d) For a synthesis of (-)-amurensinine, see: Carillo, L.; Badia, D.; Domínguez, E.; Vicario, J. L.; Tellitu, I. J. *J. Org. Chem.* **1997**, *62*, 6716. (e) Shinohara, T.; Takeda, A.; Toda, J.; Sano, T. *Heterocycles* **1998**, *48*, 981. (f) Hanessian, S.; Mauduit, M. *Angew. Chem., Int. Ed.* **2001**, *40*, 3810. (g) Dragoli, D. R.; Burdett, M. T.; Ellman, J. A. *J. Am. Chem. Soc.* **2001**, *123*, 10127.
- (13) For our preliminary communication of the enantioselective synthesis of (+)-amurensinine, see: Tambar, U. K.; Ebner, D. C.; Stoltz, B. M. *J. Am. Chem. Soc.* **2006**, *128*, 11752.
- (14) For a review of chemistry and biology of the *Lobelia* alkaloids, see: Felpin, F.-X.; Lebreton, J. *Tetrahedron* **2004**, *60*, 10127.

- (15) For syntheses of racemic (±)-lobeline, see: (a) Wieland, H.; Drishaus, I. *Annalen* **1929**, *473*, 102. (b) Wieland, H.; Koschara, W.; Dane, E. *Annalen* **1929**, *473*, 118. (c) Scheuing, G.; Winterhalder, L. *Justus Liebigs Ann. Chem.* **1929**, *473*, 126. (d) Wieland, H.; Koschara, W.; Dane, E.; Renz, J.; Schwarze, W.; Linde, W. *Justus Liebigs Ann. Chem.* **1939**, *540*, 103. (e) Parker, W.; Raphael, R. A.; Wilkinson, D. I. *J. Chem. Soc.* **1959**, 2433. (f) For an X-ray crystallography study of (-)-lobeline salts, see: Glaser, R.; Hug, P.; Drouin, M.; Michael, A. *J. Chem. Soc., Perkin Trans. 2* **1992**, 1071. (g) For a study of the biosynthesis of (-)-lobeline, see: Keogh, M. F.; O'Donovan, D. G. *J. Chem. Soc.* **1970**, 2470.
- (16) (a) Millsbaugh, C. F. *Lobelia inflata. American medicinal plants: an illustrative and descriptive guide to plants indigenous to and naturalized in the United States which are used in medicine*; Dover: New York, 1974; pp 385–388. (b) Dwoskin, L. A.; Crooks, P. A. *Biochem. Pharmacol.* **2002**, *63*, 89. (c) Thayer, A. M. *Chem. Eng. News* **2006**, *84* (39), 21.
- (17) Zheng, G.; Dwoskin, L. P.; Deacuic, A. G.; Norrholm, S. D.; Crooks, P. A. *J. Med. Chem.* **2005**, *48*, 5551.
- (18) The epimerization at C(3) is believed to be a base-catalyzed equilibration via transient retro-conjugate addition intermediate 19. See also refs 19a, 19b, and 19d.




- (19) For enantioselective syntheses of (-)-lobeline, see: (a) Compere, D.; Marazano, C.; Das, B. C. *J. Org. Chem.* **1999**, *64*, 4528. (b) Felpin, F.-X.; Lebreton, J. *J. Org. Chem.* **2002**, *67*, 9192. (c) Klingler, F.-D.; Sobotta, R. (Boehringer Ingelheim). U.S. Patent 2006014791, 2006. (d) Birman, V. B.; Jiang, H.; Li, X. *Org. Lett.* **2007**, *9*, 3237.
- (20) (a) Marion, L.; Lavigne, R.; Lemay, L. *Can. J. Chem.* **1951**, *29*, 347. (b) Franck, B. *Chem. Ber.* **1958**, *91*, 2803. (c) Logar, C.; Mesicek, M.; Perpar, M.; Seles, E. *Farm. Vestn.* **1974**, *21*; *Chem. Abstr.* **1975**, *82*, 81926h. (d) Krasnov, E. A.; Petrova, L. V.; Bekker, E. F. *Khim. Prir. Soedin.* **1977**, *585*; *Chem. Abstr.* **1977**, *87*, 164249k.
- (21) For a review of syntheses of the *Sedum* alkaloids, see: (a) Bates, R. W.; Sa-Ei, K. *Tetrahedron* **2002**, *58*, 5957. For recent syntheses of either enantiomer of sedamine, see: (b) Bates, R. W.; Nemeth, J. A.; Snell, R. H. *Synthesis* **2008**, 1033. (c) Fustero, S.; Jiménez, D.; Moscardó, J.; Catalán, S.; del Pozo, C. *Org. Lett.* **2007**, *9*, 5283. (d) Yadav, J. S.; Reddy, M. S.; Rao, P. P.; Prasad, A. R. *Synthesis* **2006**, 4005. (e) Yadav, J. S.; Reddy, M. S.; Rao, P. P.; Prasad, A. R. *Tetrahedron Lett.* **2006**, *47*, 4397. (f) Bates, R. W.; Boonsoombat, J. *Org. Biomol. Chem.* **2005**, *3*, 520. (g) Josephson, N. S.; Snapper, M. L.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2004**, *126*, 3734. (h) Zheng, G.; Dwoskin, L. P.; Crooks, P. A. *J. Org. Chem.* **2004**, *69*, 8514. (i) Angoli, M.; Barilli, A.; Lesma, G.; Passarella, D.; Riva, S.; Silvani, A.; Danieli, B. *J. Org. Chem.* **2003**, *68*, 9525. (j) Cossy, J.; Willis, C.; Bellosta, V.; BouzBouz, S. *J. Org. Chem.* **2002**, *67*, 1982.

Scheme 3



Scheme 4



3. We believed that $(-)\text{-}1$ could be accessed from amino alcohol derivative **9** via dehydration of the tertiary alcohol to form the trisubstituted olefin and subsequent removal of the sulfonamide protecting groups. Sulfonamide **9** could in turn be derived from amino-diol derivative **10** via cyclization to form the 3,4,5,6-tetrahydroazepino[5,4,3-*cd*]indole ring system of the natural product. Diol **10** could then be derived from enantioenriched diol **12** via amino alcohol derivative **11**. Diol **12** could then be accessed from the known aldehyde **13**.

Retrosynthesis of $(+)\text{-Amurensinine}$. Our approach to the isopavines, and specifically to amurensinine, is depicted in Scheme 4. Disconnection of the bridging amine in the natural product reveals azidoester **14**, which can be derived from hydroxyester $(-)\text{-}15$. We envisioned the enantioselective synthesis of this benzylic alcohol by Pd-catalyzed oxidative kinetic resolution. Hydroxyester $(\pm)\text{-}15$ could then arise from ketoester $(\pm)\text{-}16$, which we believed could be derived from arylsilyl triflate **17** and β -ketoester **18** via aryne C–C insertion²² methodology recently developed in our laboratory.

Retrosynthesis of $(-)\text{-Lobeline}$. We believed that the late-stage Pd-catalyzed oxidative desymmetrization of *meso*-diol **20**, a natural product known as lobelanidine, could deliver $(-)\text{-lobeline } ((-)\text{-}3$, Scheme 5). We also envisioned that the diastereoselective formation of the desired *cis*-2,6-disubstituted piperidine moiety of amino alcohol $(-)\text{-}3$ could be achieved

under equilibrating conditions via intermediate enone **19**. Diol **20** could then be derived via disconnection of a C–C bond to arrive at amino alcohol $(\pm)\text{-}21$. Enantioselective oxidation of amino alcohol derivative $(\pm)\text{-}21$ in turn would allow us to access either $(-)\text{-sedamine } ((-)\text{-}4)$ or its enantiomer. We believed that amino alcohol derivative $(\pm)\text{-}21$ could be readily accessed from aldehyde **22**.

Synthesis of $(-)\text{-Aurantoclavine}$. Our synthesis of the ergot alkaloid $(-)\text{-aurantoclavine}$ commenced with aldehyde **13** (Scheme 6).²³ Addition of the dianion derived from isobutylene oxide²⁴ to the aldehyde furnished racemic diol $(\pm)\text{-}12$. At this stage, application of a Pd-catalyzed oxidative kinetic resolution delivered enantioenriched diol $(-)\text{-}12$ in 96% ee and 37% yield (91% of the theoretical maximum, selectivity factor²⁵ $s = 18.2$). Product ketone **23** could be readily recycled by reduction with lithium aluminum hydride, affording diol $(\pm)\text{-}12$ in 95% yield.


The enantioenriched diol $(-)\text{-}12$ was then transformed to tricyclic alcohol **9** (Scheme 7). Diol $(-)\text{-}12$ was treated with hydrazoic acid under Mitsunobu conditions²⁶ to furnish azidoalcohol **24**. This transformation was conducted at low temperature to minimize racemization at the sensitive benzylic stereocenter. Hydrogenation of the azide and protection of the amine as a 2-nitrobenzenesulfonamide²⁷ delivered sulfonamide **11**. Bromination of the indole nucleus was followed by

(22) (a) Tambar, U. K.; Stoltz, B. M. *J. Am. Chem. Soc.* **2005**, *127*, 5340. Others have disclosed similar aryne insertions since our initial report, see: (b) Yoshida, H.; Watanabe, M.; Ohshita, J.; Kunai, A. *Chem. Commun.* **2005**, 3292. (c) Yoshida, H.; Watanabe, M.; Ohshita, J.; Kunai, A. *Tetrahedron Lett.* **2005**, *46*, 6729.


(23) Kozikowski, A. P.; Ishida, H.; Chen, Y.-Y. *J. Org. Chem.* **1980**, *45*, 3350.

(24) Bachki, A.; Foubelo, F.; Yus, M. *Tetrahedron: Asymmetry* **1996**, *7*, 2997.


Scheme 5



Scheme 6



Scheme 7




vinylation of bromoindole **25** under Stille conditions²⁸ to furnish vinyl indole **26**. Hydroboration–oxidation²⁹ of the olefin of **26** to amino diol derivative **10** was followed by formation of the azepine under Mitsunobu conditions to deliver tricyclic alcohol


9 in excellent yield. We then investigated the dehydration of the tertiary alcohol **9** to (-)-aurantioclavine (Scheme 8).

Unfortunately, under various reaction conditions,³⁰ dehydration of the tertiary alcohol delivered a mixture of inseparable

Scheme 8



Scheme 9




olefin isomers, with the undesired isomer **28** predominating in all cases. In an attempt to access the desired trisubstituted olefin isomer, we then turned to dehydration of intermediate **26** prior to formation of the azepine ring (Scheme 8).

We found that employing phosphorus oxychloride as the dehydrating agent in pyridine as solvent delivered the desired trisubstituted olefin isomer **29** as the major product. The olefin isomers **29** and **30** were separable by employing silica gel impregnated with silver nitrate.³¹

The completion of the synthesis of (-)-aurantioclavine is depicted in Scheme 9. Selective hydroboration–oxidation of the terminal olefin in vinyl indole **29** delivered the protected amino alcohol **31**, which was then transformed to the tricyclic alcohol **27** under Mitsunobu conditions in excellent yield. Removal of the *o*-nitrobenzenesulfonyl protecting group delivered amine **32**. Subsequent removal of the tosyl group using tetrabutylammonium fluoride (TBAF)³² then delivered (-)-aurantioclavine ((-)-**1**).

Scheme 10



Synthesis of (+)-Amurensinine. The synthesis of amurensinine began from diazoketone **33**, readily available in five steps and 95% overall yield from homoveratric acid (Scheme 10). Selective intramolecular C–H insertion provided β -ketoester **18** in excellent yield.³³ Treatment of β -ketoester **18** with arylsilyl

(25) The selectivity factor, s , for kinetic resolution is defined as the ratio of the rate of reaction of the fast-reacting enantiomer of the racemic substrate, k_{fast} , to the rate of reaction of the slow-reacting enantiomer, k_{slow} , in the same transformation, i.e., $s = (k_{\text{fast}}/k_{\text{slow}})$. The selectivity factor can be expressed in terms of enantiomeric excess (ee) of the remaining alcohol and the conversion (c) of the alcohol to the corresponding ketone, i.e., $s = \ln[(1-c)(1-ee)]/\ln[(1-c)(1+ee)]$. For $s > 10$, synthetically useful quantities of enantioenriched alcohol can be accessed. For example, an oxidative kinetic resolution with $s = 10$ at 62% conversion would afford recovery of alcohol of 90% ee. See also: Kagan, H. B.; Fiaud, J. C. In *Topics in Stereochemistry*; Eliel, E. L., Ed.; Wiley & Sons: New York, 1988; Vol. 18, pp 249–330.

(26) Mitsunobu, O. *Synthesis* **1981**, 1.

(27) Fukuyama, T.; Jow, C.-K.; Cheung, M. *Tetrahedron Lett.* **1995**, 36, 6373.


(28) For reviews of the Stille cross-coupling reaction, see: (a) Stille, J. K. *Angew. Chem., Int. Ed. Engl.* **1986**, 25, 508. (b) Farina, V.; Krishnamurthy, V.; Scott, W. J. *Org. React.* **1997**, 50, 1. (c) Littke, A. F.; Fu, G. C. *Angew. Chem., Int. Ed.* **2002**, 41, 4176. (d) Espinet, P.; Echavarren, A. M. *Angew. Chem., Int. Ed.* **2004**, 43, 4704.

(29) (a) Knights, E. F.; Brown, H. C. *J. Am. Chem. Soc.* **1968**, 90, 5281. (b) Brown, H. C.; Knights, E. F.; Scouten, C. G. *J. Am. Chem. Soc.* **1974**, 96, 7765.


(30) See Supporting Information for details.

(31) Morris, L. J. *J. Lipid Res.* **1966**, 7, 717.


Scheme 11



Scheme 12



Scheme 13




triflate **17**, available in five steps from sesamol, in the presence of cesium fluoride^{22a,34} afforded ketoester (\pm)-**16**, thereby generating the polycyclic carbon framework of amurensinine in rapid fashion.


Chemo- and diastereoselective reduction of the ketone of (\pm)-**16** with L-Selectride generated hydroxyester (\pm)-**15** as a single diastereomer, presumably resulting from equatorial delivery of hydride to the ketone (Scheme 11).

We then examined the Pd-catalyzed oxidative kinetic resolution of hydroxyester (\pm)-**15**. Under optimized reaction conditions, the racemic alcohol could be resolved to highly enantioenriched hydroxyester (-)-**15** with good selectivity (Scheme

Scheme 14



Scheme 15




12). Though the overall mass recovery of (-)-**15** and (+)-**16** was not ideal, significant quantities of enantioenriched alcohol (-)-**15** could be accessed from this pathway.

We next investigated introduction of the amine required for amurensinine from alcohol (-)-**15**. We found that alcohol (-)-**15** was unreactive to many common nitrogen-bearing nucleophiles under Mitsunobu conditions, or it reacted to form a stilbene system by elimination. However, we were able to install an azide by employing diphenylphosphoryl azide in a procedure developed specifically for electron-rich benzylic alcohols by Thompson and co-workers (Scheme 13).³⁵ Reduction of the resulting azide afforded bridged lactam (+)-**34**. Lactam reduction and reductive methylation afforded (+)-amurensinine ((+)-**2**).

(35) Thompson, A. S.; Humphrey, G. R.; DeMarco, A. M.; Mathre, D. J.; Grabowski, E. J. J. *J. Org. Chem.* **1982**, 47, 4808.

- (32) Yasuhara, A.; Sakamoto, T. *Tetrahedron Lett.* **1998**, 39, 595.
(33) (a) Wenkert, E.; Davis, L. L.; Mylari, B. L.; Solomon, M. F.; Da Silva, R. R.; Shulman, S.; Warnet, R. J.; Ceccherelli, D.; Curini, M.; Pellicciari, R. J. *J. Org. Chem.* **1982**, 47, 3242. (b) Taber, D. F.; Petty, E. H. *J. Org. Chem.* **1982**, 47, 4808.
(34) Himeshima, Y.; Sonoda, T.; Kobayashi, H. *Chem. Lett.* **1983**, 1211.


Scheme 16



While (+)-amurensinine could be accessed by this route, the sequence to (+)-2 from (±)-15 was complicated by the formation of several side products. Treatment of hydroxyester (–)-15 with DBU in the absence of diphenylphosphoryl azide produced lactone 35, demonstrating the sensitivity of the bis-benzylic stereocenter to epimerization (Scheme 14). Of further concern was the formation of lactam (+)-34 in only 57% ee. This partial racemization required two stereocenters to be inverted in the azide installation reaction, presumably via an intermediate such as *o*-quinonodimethide 36.

Our difficulty in suppressing epimerization at the bis-benzylic stereocenter bearing the ester in the azide displacement led us to pursue an alternate route to amurensinine via intermediates with attenuated acidity at the bis-benzylic carbon (Scheme 15). Also, we hoped an alternate benzylic alcohol substrate would lead to improved mass recovery in the oxidative kinetic resolution. To this end, reduction of hydroxyester (±)-15 was followed by selective silylation of the primary alcohol to afford alcohol (±)-37. Oxidative kinetic resolution of this alcohol proved highly selective (selectivity factor $s > 47$), providing highly enantioenriched alcohol (–)-37. Interestingly, resolutions that were allowed to proceed to high ee of (–)-37 did not afford any of the expected ketone (+)-39. Instead, diketone (–)-38 was formed in good yield and high ee. Monitoring of the reaction demonstrated that the ketone (+)-39 was being generated, but it slowly underwent further oxidation to the diketone (–)-38. In fact, isolated samples of ketone 39 were slowly oxidized to the diketone in C₆D₆. Handling of this ketone under


Scheme 18



^a Isolated yield. ^b Measured by chiral HPLC after derivatization.


^c Equivalents of phenoxide relative to Pd(sparteine)Cl₂. The phenoxide was stirred with Pd(sparteine)Cl₂ for 1 h prior to introduction of substrate.

Scheme 19




argon delayed this oxidative decomposition. We hypothesized that ketone (+)-39 was reacting with molecular oxygen via a radical pathway to afford the diketone (–)-38. This hypothesis is supported by our observations on non-enantioselective oxidations performed on alcohol (±)-37. Thus, oxidation of (±)-37 with Dess–Martin periodinane³⁶ cleanly provided ketone (±)-39, while diketone (±)-38 was obtained when either MnO₂ or Pd(OAc)₂/O₂³⁷ was employed as the oxidant. We therefore screened various radical inhibitors as additives in kinetic resolutions of alcohol (±)-37. Tetracyanoethylene led to little alcohol oxidation. Neither BHT nor 2-methyl-2-butene suppressed ketone overoxidation; however, incorporation of even catalytic quantities of 2-methyl-2-butene enhanced mass recovery and improved selectivity in the kinetic resolution. While the role of 2-methyl-2-butene is not yet clear, it was included in our preparative experiments because of its beneficial effect. Thus, under our optimized conditions, (±)-37 was resolved with Pd(sparteine)Cl₂ as a catalyst in the presence of O₂ to 99% ee and was isolated in 47% yield (94% of the theoretical maximum).

Scheme 17



Scheme 20



Having accessed substantial quantities of highly enantioenriched alcohol $(-)$ -**37**, we next proceeded with the installation of the nitrogen required in the natural product. Azide displacement under Thompson's conditions was followed by desilylation to afford azidoalcohol $(-)$ -**40** (Scheme 16). Gratifyingly, this azide was obtained with clean inversion in 99% ee, indicating that decreasing the acidity of the bis-benzylic center did indeed minimize $\text{S}_{\text{N}}1$ pathways. Oxidation of alcohol $(-)$ -**40** in two steps to the corresponding carboxylic acid and subsequent azide reduction delivered lactam $(+)$ -**34**. Reduction and methylation as before afforded $(+)$ -amurensinine $(+)$ -**2** in 99% ee.³⁸

Synthesis of $(-)$ -Lobeline and $(-)$ - and $(+)$ -Sedamine. Our synthesis commenced from known aldehyde **22**.³⁹ Addition of the anion derived from Horner–Emmons reagent **41** followed by hydrolysis of the resulting methyl enol ether provided ketone (\pm) -**42** in good overall yield (Scheme 17). DIBAL emerged as the preferred reductant from a screen of conditions for the subsequent reduction of aryl ketone (\pm) -**42**, providing the amino alcohol derivative (\pm) -**21** with 12:1 diastereoselectivity.⁴⁰ The diastereomers were separable by chromatography, and the benzylic alcohol of the major diastereomer (\pm) -**21** was protected as a TBS ether. Formylation with DMF under anionic conditions furnished aldehyde **43** as a single diastereomer. Aldehyde **43** was then treated with the anion derived from Horner–Emmons reagent **41**. Hydrolysis of the intermediate enol ether afforded ketone **44**. As before, diastereoselective ketone reduction was accomplished with DIBAL, providing a single alcohol diastereomer. Subsequent desilylation with TBAF followed by exhaustive reduction of the carbamate delivered the *meso*-diol lobelanidine (**20**). The relative stereochemistry of lobelanidine was confirmed by X-ray crystallography.

At this stage, we investigated the key desymmetrization of **20** to access $(-)$ -lobeline $(-)$ -**3**. Our previously optimized reaction conditions,⁴¹ involving catalytic Pd(sparteine)Cl_2 in chloroform under an oxygen atmosphere, furnished lobeline (**3**) as a 1:1 mixture of diastereomers at C(3) in 70% yield and 95% ee (Scheme 18).⁴² Synthetic alkaloid **3** was found to be spectroscopically identical to the equilibrated natural isolate and possessed identical optical rotation. Interestingly, a survey of additives revealed that incubation of Pd(sparteine)Cl_2 with the sodium salt of 6-methoxy-2-naphthol (**45**) for 1 h prior to exposure to *meso*-diol **20** had a dramatic impact on the reaction. Inclusion of the phenoxide salt permitted smooth conversion to amino alcohol $(-)$ -**3** with a lowered catalyst loading (10 mol %) and reaction time (48 h) and furnished amino alcohol $(-)$ -**3** in enhanced isolated yield (87%) and ee (99%). Although the role of the phenoxide is presently unclear, one possibility is that the observed improvement in overall reaction profile may result from the formation of a more reactive, catalytically competent $\text{Pd(sparteine)phenoxide}$ salt.

After implementing a late-stage Pd-catalyzed oxidative desymmetrization to access $(-)$ -lobeline $(-)$ -**3** as a 1:1 mixture

of equilibrating diastereomers, we began efforts to control the epimerization at C(3) in order to access $(-)$ -**3** as a single diastereomer. The rate of epimerization of natural $(-)$ -**3** was found to increase with increasing basicity of the reaction medium and by protic solvents.⁴³ Furthermore, we found that the hydrochloride salt **3**·HCl (*cis:trans* = 1:1) could be equilibrated to a 3:1 mixture of the *cis:trans* isomers in *i*-PrOH. Prompted by these observations, we developed a dynamic crystallization method allowing for selective precipitation of the *cis*-isomer of aminoketone **3** (Scheme 19). Heating a 1:1 mixture of *cis:trans* isomers of the hydrochloride of aminoketone **3** in *i*-PrOH followed by slow crystallization permitted a 69% recovery of *cis*- $(-)$ -**3** after liberation of the free base.

Having demonstrated a concise total synthesis of isomerically pure $(-)$ -lobeline $(-)$ -**3**, we explored the utility of protected amino alcohol (\pm) -**21** for the synthesis of related *Sedum* alkaloids, namely, both enantiomers of sedamine (i.e., $(-)$ - and $(+)$ -**4**, Scheme 20). Exposure of protected amino alcohol (\pm) -**21** to our optimized Pd(II)-catalyzed aerobic oxidative kinetic resolution conditions furnished protected amino alcohol $(-)$ -**21** in 94% ee and the oxidation product, ketone $(-)$ -**42**, in 81% ee with excellent yield and selectivity.

Direct reduction of amino alcohol derivative $(-)$ -**21** with lithium aluminum hydride provided natural $(-)$ -sedamine $(-)$ -**4**. Reiteration of the highly diastereoselective DIBAL reduction of ketone $(-)$ -**42** generated enantiomeric alcohol $(+)$ -**21**, and further reduction of the carbamate delivered the non-natural $(+)$ -sedamine $(+)$ -**4**.

Conclusion

Herein, we have demonstrated the power of palladium-catalyzed enantioselective aerobic oxidation toward the synthesis of alkaloid natural products. We have developed an enantioselective synthesis of the ergot alkaloid $(-)$ -aurantioclavine, a promising intermediate en route to the communesin family of alkaloids. We have also demonstrated that the strategic com-

(36) Dess, D. B.; Martin, J. C. *J. Org. Chem.* **1983**, *48*, 4155.

(37) Nishimura, T.; Onoue, T.; Ohe, K.; Uemura, S. *J. Org. Chem.* **1999**, *64*, 6750.

(38) We have recently accomplished a formal synthesis of the naturally occurring enantiomer of amurensinine using a readily accessible diamine ligand that serves as a mimic for $(+)$ -sparteine in the oxidative kinetic resolution. See: Ebner, D. C.; Trend, R. M.; Genet, C.; McGrath, M. J.; O'Brien, P.; Stoltz, B. M. *Angew. Chem., Int. Ed.* **2008**, *47*, 6367–6370.

(39) Molander, G. A.; Romero, J. A. C. *Tetrahedron* **2005**, *61*, 2631.

(40) A rationale for the observed selectivity follows from the calculated conformational equilibria of *cis,cis*-1,4-cyclooctadiene. See: Anet, F. A. L.; Yavari, I. *J. Am. Chem. Soc.* **1977**, *99*, 6986. Also see Supporting Information for a stereochemical model.

(41) Bagdanoff, J. T.; Stoltz, B. M. *Angew. Chem., Int. Ed.* **2004**, *43*, 353.

(42) The enantiomeric excess was determined after derivatization of lobeline by the method described in ref 21h. See Supporting Information for details.

bination of an aryne insertion followed by enantioselective oxidation allows efficient access to the isopavine alkaloids, exemplified by (+)-amurensinine. Finally, our concise enantioselective synthesis of (–)-lobeline in diastereomerically pure form, as well as the synthesis of either enantiomer of sedamine, demonstrates the extraordinarily facile generation of multiple stereogenic centers in a catalytic enantioselective manner in a single operation.

Acknowledgment. The authors thank the NIH-NIGMS (R01 GM65961-01), California TRDRP (postdoctoral fellowships to Y.K.R. and S.K., and predoctoral fellowship to J.T.B.), NDSEG (predoctoral fellowships to D.C.E. and U.K.T.), NSF (predoctoral fellowship to D.C.E.), Caltech, AstraZeneca, Boehringer Ingelheim,

Johnson & Johnson, Pfizer, Merck, Amgen, Abbott, Research Corporation, Roche, and GlaxoSmithKline for generous funding.

Supporting Information Available: Full experimental details and characterization data for all new compounds (PDF, CIF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

JA804738B

(43) Crooks and co-workers^{21h} noted the rate of epimerization of diastereomerically pure *cis*-lobeline to 46:54 *cis:trans*-lobeline in various deuterated solvents by ¹H NMR (CD₃OD > CD₃CN > CD₃COCD₃ > CDCl₃). In our hands, the epimerization of diastereomerically pure *cis*-lobeline to 1:1 *cis:trans*-lobeline was much slower in C₆D₆ (5 days) as opposed to CDCl₃ (2 days) and CD₃OD (2 h).